

## SUPPLEMENTAL LABELING

# STEALTH HERBICIDE

FOR DISTRIBUTION AND USE ONLY WITHIN IDAHO  
EPA Reg. No. 34704-868      EPA SLN No.: ID- 060017

## FOR WEED CONTROL IN CLOVER GROWN FOR SEED PRODUCTION

### DIRECTIONS FOR USE

This label expires. December 31, 2019

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered product label for STEALTH herbicide, EPA Reg. No. 34704-868.

This labeling must be in the possession of the user at the time of application.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STEALTH** controls most annual grasses and certain broadleaf weeds as they germinate, but it will not control established weeds. Unusually cold, excessively wet, or hot and dry conditions that delay germination or extend it over a long period of time may reduce weed control.

**Some stunting and chlorosis of the clover may occur with postemergence applications. Applications when clover exceeds 6 inches in height may result in poor weed control in the crop row due to possible reduced spray coverage to the soil. DO NOT** exceed 4.8 quarts of product per acre in any one crop season. **DO NOT** apply this product through any type of irrigation system. Uniformly apply in 10 or more gallons of water per acre with ground equipment.

Apply **STEALTH** at optimum application timing for dodder or other weed control to established clover at a broadcast rate of 1.2 to 4.8 quarts per acre prior to weed emergence. Apply **STEALTH** before clover exceeds 6 inches in height.

**STEALTH** is most effective when mechanically incorporated, or when incorporated in the weed germination zone by rainfall or irrigation within 7 days of application.

**DO NOT** apply to seedling clover. **DO NOT** use on peat or muck soils.

**SPECIAL CROP USE RESTRICTIONS: Use of this product according to this label is deemed a non-food use and as such is not for use on field producing clover for livestock feed.**

**DO NOT** feed or graze clover following **STEALTH** application.

**DO NOT** cut treated clover for hay or forage.

**DO NOT** use harvested seed for sprouting.

No portion of the treated field, including seed, seed screenings, hay, forage, or stubble may be used for human or animal feed.

Producers of clover seed who use this product, or cause the product to be used on a field they operate, are required to inform, in writing, conditioners receiving seed produced on fields treated with STEALTH Herbicide. A copy of the label must be given by the producer to the conditioner. Processed seed must be labeled “**Not for human or animal consumption**” at the processing plant. All clover seed screenings must be disposed of in such a way that they cannot be distributed or used for food or feed.

**NOTE:** If loss of clover stand occurs due to adverse weather conditions, any crop registered for **STEALTH** preplant incorporated use can be replanted the same year into **STEALTH** treated soil without adverse effects. If replanting is necessary, soil should not be reworked deeper than the **STEALTH** treated zone.

**STEALTH herbicide** applied at a rate of 1.2 to 2.4 quarts per acre prior to weed emergence will control the following weeds:

**Grasses:** Barnyardgrass, Crabgrass, Crowfootgrass, Foxtail, Giant, Foxtail, Green, Foxtail, Yellow, Goosegrass, Johnsongrass (seedling), Panicum, Fall; Panicum, Texas; Sandbur, Field; Signalgrass, Witchgrass

**Broadleaf Weeds:** Carpetweed, Florida Pusley, Kochia, Lambsquarters, Pigweed, Purslane, Spurge, annual

**STEALTH** applied at a rate of 2.4 to 4.8 quarts per acre prior to weed emergence will control the following weeds in addition to those listed above:

**Grasses:** Bluegrass (annual), Cupgrass, Woolly; Itchgrass, Junglerice, Lovegrass, Millet, Wild-proso, Panicum, Browntop, Shattercane, Sprangletop, Mexican, Sprangletop, Red

**Broadleaf Weeds:** Chickweed (common), Dodder\*, Fiddleneck, Henbit, Knotweed (prostrate), Puncturevine, Rocket, London; Shepherdspurse, Smartweed, Pennsylvania; Velvetleaf (Buttonweed)

**ROTATIONAL CROP RESTRICTIONS**

The following crops may be planted after the indicated time interval depending on the crop and rate used.

Crop	Rate per Acre		
	Less than 1.8 quarts	1.9 to 3.6 quarts	3.7 to 4.8 quarts
Crops labeled for Preplant Incorporated Application	Same season	4 months	4 months
Wheat	4 months	12 months	18 months
Red beets, sugar beets and spinach	12 months	18 months	24 months
Other crops*	Following year	18 months	24 months

\*Barley may be planted 12 months after an application of up to 2.4 quarts per acre.

**To avoid crop injury, land should be plowed using a moldboard plow when greater than 1.8 quarts per acre are used, or when rotating to sugar beets, red beets, or spinach at any use rate.**

**24c Registrant:  
Loveland Products Inc.  
PO Box 1286  
Greeley, CO 80632-1286**