



SPECIAL LOCAL NEED 24(C) LABELING

Lorox® DF

Agricultural Herbicide

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO
EPA Reg. No. 61842-23 EPA SLN No. ID-140001**

This SLN expires and must not be used or distributed after December 31, 2015

CROP ROTATION DIRECTIONS FOLLOWING LOROX DF USE ON LENTILS, DRY PEAS AND CHICKPEAS

Directions for Use: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA registered label. This labeling must be in the possession of the user at the time of pesticide application.

Crop Rotation Directions (Following LOROX® DF use on Lentils, Dry Peas and Chickpeas):

Apply LOROX DF at a single application at up to 2 pounds per acre post planting but preemergence to the crop. Sufficient rainfall to move the herbicide into the root zone of germinating weeds is needed for best performance. Use lower rates on coarse soils and higher rates on heavier soils. LOROX DF may be tank mixed with other herbicides registered for use in these crops. When tank mixing, use the most restrictive label limitations for each of the products being used in the tank mix.

Any crop registered for the rate of LOROX DF that was applied may be replanted immediately. Winter Wheat and Winter Barley may be planted 4 months after application. After 12 months, any other crop used for food or feed may be rotated.

RESTRICTIONS/PRECAUTIONS

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
- Do not apply more than one (1) application per acre per season.
- Do not apply by air.
- Do not apply to sand or loamy sand soil types.
- Do not use on soils with less than 1% organic matter.
- Do not apply this product to dry peas through any type of irrigation system.
- Do not apply more than 1.25 pounds per acre of LOROX DF (0.625 pounds of active ingredient) per acre per year, if the treated lentil field will be rotated into winter wheat in the fall or winter and where annual rainfall average is less than 16 inches.
- Do not plant spring wheat, barley, canola or any other crops not registered for the rate of linuron that was applied until 12 months after the last application of LOROX DF.

For product information, call 1-800-525-2803. You may also contact 1-866-374-1975 for emergency medical treatment information.

This Section 24(c) SLN label contains new or supplemental instructions for use of this product which

do not appear on the EPA-registered label. Follow the instructions carefully.

To the extent consistent with applicable law, Tessenderlo Kerley, Inc makes this product available to the user and/or grower solely to the extent the benefit and utility, in the sole opinion of the user and/or grower, outweigh the extent of potential injury associated with the use of this product. The decision to use or not to use LOROX DF must be made by each individual LOROX DF user and/or grower on the basis of possible crop injury from LOROX DF, the severity of weed infestation, the cost of alternative weed controls, and other factors. Read the Warranty and Limitation of Damages on the product label before buying or using this product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Warranty and Limitation of Damages on the product label.

Section 24(c)

Rev. 04/01/2014

Tessenderlo Kerley, Inc.
2255 N. 44th Street, Suite 300 Lorox® and NovaSource® are registered trademarks of Tessenderlo Kerley, Inc
Phoenix, AZ 85008 USA ©2013, 2014 Tessenderlo Kerley, Inc. All Rights Reserved
1-800-525-2803
www.novasource.com