

Special Local Need (SLN) Label



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Opensight[®]
EPA Reg. No. 62719-597

EPA 24(c) Special Local Need Registration SLN ID-090018
For Distribution and Use Only Within the State of Idaho

This label for Opensight expires and must not be distributed or used in accordance with this SLN registration after December 31, 2019.

For Control of Herbaceous Broadleaf Weeds and Woody Plants in Forest Plantations and Grazed Areas In and Around These Sites

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read this SLN labeling and the label affixed to the container for Opensight[®] specialty herbicide before applying. All applicable use directions, precautions and restrictions on this SLN labeling and the label affixed to the container must be followed.

Directions for Use

Refer to product package label for further use directions including requirements for the Worker Protection Standard (WPS), Use Precautions and Restrictions, Precautions for Avoiding Spray Drift, Mixing Instructions, Application Directions, and Broadleaf Weeds Controlled.

Forest Plantations

Opensight[®] specialty herbicide may be applied to forest plantations, as an aerial or ground broadcast application for site preparation, or as, a spot application or directed spray application for conifer release (see Application Methods section) to control herbaceous broadleaf weeds and woody plants. Opensight may be applied alone or in tank-mix combinations with labeled rates of other herbicides provided: (1) the tank mix product is labeled for the timing and method of application for the use site to be treated and (2) mixing is not prohibited by the label of the registered tank mixed products. Use as directed in the Directions of Use section of the tank-mix partner.

Follow Mixing Instructions under the General Mixing and Application Instructions section below. Unless otherwise directed, use an agricultural labeled crop oil concentrate, methylated seed oil, or non-ionic surfactant for all treatments. Apply surfactants at a minimum rate of 0.25% v/v (1 quart per 100 gallons spray solution) or at the manufacturers recommended rate.

Application Methods

For best control from broadcast and directed spray applications of Opensight, use a spray volume which will provide thorough plant coverage. Recommended spray volumes are usually 10 to 30 gallons per acre by air or 10 to 100 gallons per acre by ground.

Forest Site Preparation (Not for Conifer Release)

Use up to 3.3 ozs of Opensight and apply in a total spray volume of 10 to 30 gallons per acre. Tank mixtures with other herbicides registered for forest use, such as Accord XRT II, may improve control of woody brush, grasses, and herbaceous plants. When tank mixtures of herbicides are used for forest site preparation, labels for all products in the mixture must be followed; use the most restrictive label directions, including the longest recommended waiting period before planting.

Note: Conifers planted sooner than 4 months after treatment with Opensight may be injured.

Directed Spray Applications for Conifer Release

NOTE: To prevent injury to or loss of desirable trees, including conifers, avoid spray contact with the foliage, especially small trees and their terminal buds, and in the root zone when making directed spray applications or spot treatments.

Use Opensight to release conifers from competing invasive and broadleaf weeds such as common crupina, hawkweed, knapweeds, St. Johnswort, thistles, and sensitive woody plants such as blackberry, buckbrush, snowberry, locust, redbud, and wild roses. The spray mixture should be directed onto foliage of competitive plants using knapsack or backpack sprayers with flat fan nozzles or equivalent any time after plants have reached full leaf size, but before autumn coloration. The majority of treated hardwoods should be less than 6 feet in height to ensure adequate spray coverage. Care should be taken to direct spray away from contact with conifer foliage, particularly foliage of desirable conifers.

Spot Application for site preparation or conifer release: Spot applications may be made at rates equivalent to the broadcast-applied rates of 2 to 3.3 oz. per acre per annual growing season depending on the target species. Spray volume should be sufficient to thoroughly and uniformly wet weed foliage. Repeat treatments may be made, but the total amount of Opensight applied must not exceed 3.3 oz per acre per year. To prevent misapplication, spot treatments should be applied using calibrated application techniques. Apply the spray mixture using a calibrated boom, boomless spray system, hand-held, or backpack sprayers.

Spot applications may be made at a rate of up to 0.22 lb acid equivalent aminopyralid (6.6 oz of Opensight) per acre; however, not more than 50% of an acre may be treated. Do not apply more than a total of 0.11 lb acid equivalent aminopyralid per acre (3.3 oz per acre of Opensight) per annual growing season as a result of broadcast, spot or repeat applications.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow
R372-015
Accepted: __/__/__
Replaces: R372-001