



**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO**

**EVEREST® 2.0 HERBICIDE**

**EPA Reg. No. 66330-391**

**EPA SLN No. ID-110003**

**For Application on Kentucky Bluegrass Grown For Seed (Only During the Year of Establishment) in the State of Idaho  
FIFRA §24(c) Special Local Need Registration**

**Active Ingredient**

**Flucarbazono-sodium,**

**4,5-Dihydro-3-methoxy-4-methyl-5-oxo-N-**

**[[2-(trifluoromethoxy)phenyl]sulfonyl]-1H-**

**1,2,4-triazole-1-carboxamide, sodium salt..... 35.0%**

**Other Ingredients ..... 65.0%**

**Total..... 100.0%**

**This formulation contains 3.5 lbs of Flucarbazono-Sodium active ingredient per gallon (419 g ai/l)**

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

**In Case of Medical Emergency Involving This Product,  
Call Day or Night: 800-228-5635, Ext. 174**

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling and the main label must be in the possession of the user at the time of application.

Read the entire label for EVEREST 2.0 HERBICIDE before proceeding with the use directions contained in this supplemental labeling. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label and this supplemental label.

## ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water when disposing of equipment washwaters or rinsate.

Do not allow sprays to drift onto adjacent desirable plants.

## USE RATES

Apply EVEREST 2.0 Herbicide to Kentucky bluegrass grown for seed (only during the year of establishment) for control of wild oats, windgrass, and other grass and broadleaf weeds at a rate of 1.0 fl oz/A, with a non-ionic surfactant at a rate of 0.25% v/v.

Recommended Rates for Kentucky Bluegrass Grown For Seed		
Grass Weeds	Rate (fl oz/A)	Notes
Wild Oats ( <i>Avena fatua</i> )	1.0 oz/A	Control
Windgrass ( <i>Apera spica-venti</i> )	1.0 oz/A	Control
Cheat ( <i>Bromus secalinus</i> )	1.0 oz/A	Control with fall applications. Suppression with spring applications.
Ventenata ( <i>Ventenata dubia</i> )	1.0 oz/A	Suppression
Broadleaf Weeds	Rate	Notes
Wild Mustard	1.0	Control
Black Mustard	1.0	Control
Blue Mustard	1.0	Control
Tansy Mustard	1.0	Control
Tumble Mustard	1.0	Control

## GROUND APPLICATION

Apply in water using a spray volume of 5 to 10 gallons/acre (or 50 to 100 liters/hectare) at 30 to 50 psi to ensure proper weed coverage. Flat fan nozzles of 80 or 110 degrees are recommended for optimum coverage. Do not use floodjet or controlled droplet application (CDA) equipment. Nozzles may be oriented 45 degrees forward to enhance crop penetration and to give better weed coverage.

## AERIAL APPLICATION

Apply in water using a minimum spray volume of 3 gal/A (or 30 L/ha). For best results, particularly in dry conditions or where heavy weed infestations exist, use a minimum of 5 gal/A (or 50 L/ha). Use nozzles that provide 200 to 350 micron size droplets for best results and to insure uniform spray coverage. Aerial applications with EVEREST 2.0 Herbicide should be made with low drift nozzles at a maximum height of 10 feet above the crop and at a maximum pressure of 40 psi. Do not apply aurally when wind speed is greater than 10 mph. Do not allow spray to drift onto adjacent crops, as injury or loss may occur.

Refer to the **SPRAY DRIFT MANAGEMENT** section of the federal label for additional information on how to reduce drift during aerial application.

## ENDANGERED SPECIES PROTECTION

To avoid adverse effects on endangered dicot plant species, the following measures will be required where endangered plant species occur in the counties listed in the table below:

State	County
Idaho	Benewah Clearwater Idaho Kootenai Latah Lewis Nez Perce

For ground applications, the applicator must:

- Apply when there is sustained wind away from native plant communities, OR
- Use low-pressure nozzles according to manufacturer's specifications that produce only coarse or very coarse droplets, OR
- Leave a 50 foot untreated buffer between the treatment and native plant communities.

For aerial applications, the applicator must:

- Apply only when there is sustained wind away from native plant communities, OR
- Leave a 350 foot untreated buffer between the treatment and native plant communities.

## RESTRICTIONS/PRECAUTIONS

- Use of this product according to this labeling is deemed a non-food use by the Idaho Department of Agriculture, and as such is not for use on fields producing bluegrass for livestock feed.
- Do not feed or graze bluegrass following application of EVEREST 2.0 Herbicide.
- Do not cut treated bluegrass for hay, or for forage.
- Do not use harvested seed for sprouting.
- No portion of the treated field, including seed, seed screenings, hay, forage or stubble may be used for human or animal feed.
- Producers of bluegrass seed who use this product, or cause the product to be used on a field that they operate are required to inform, in writing, conditioners receiving seed produced on fields treated with EVEREST 2.0 Herbicide.
- A copy of the labeling must be given by the producer to the conditioner.
- Processed seed must be labeled "**Not for human or animal consumption**".
- All bluegrass seed screenings, hay, forage and other crop by-products produced from treated fields must be disposed of in such a way that they cannot be distributed or used for food or feed.
- Do not apply more than 1.0 fl oz/A of EVEREST 2.0 Herbicide per year.
- Do not make more than one postemergence application of EVEREST 2.0 Herbicide per year.

- Do not use EVEREST 2.0 Herbicide in annual production systems of Kentucky bluegrass grown for seed.
- For use under this SLN label do not apply EVEREST 2.0 Herbicide through any type of irrigation system.
- Do not apply EVEREST 2.0 Herbicide within 1 year (365 days) of harvesting Kentucky bluegrass seed.
- This product is for use during the establishment year only. Do not graze livestock or harvest forage from treated Kentucky bluegrass seed fields within 1 year (365 days) of application.

**ARYSTA LIFESCIENCE NORTH AMERICA, LLC'S WARRANTY DISCLAIMER FOR USE OF EVEREST 2.0 HERBICIDE ON KENTUCKY BLUEGRASS GROWN FOR SEED**

EVEREST 2.0 Herbicide when used on Kentucky bluegrass grown for seed may lead to crop injury, loss, or damage. Arysta LifeScience North America, LLC recommends that the user and/or grower test EVEREST 2.0 Herbicide in order to determine its suitability for such intended use. Arysta LifeScience North America, LLC makes EVEREST 2.0 Herbicide available to the user and/or grower solely to the extent the benefit and utility, in the sole opinion of the user and/or grower, outweigh the extent of potential injury associated with the use of this product. The decision to use or not to use EVEREST 2.0 Herbicide must be made by each individual EVEREST 2.0 Herbicide user and/or grower on the basis of possible crop injury from EVEREST 2.0 Herbicide, the severity of weed infestation, the cost of alternative weed controls, and other factors. Arysta LifeScience North America, LLC intends that because of the risk of crop damage that all such use is at the user's and/or grower's risk. Arysta LifeScience North America, LLC makes no warranty, express or implied, with respect to the suitability of EVEREST 2.0 Herbicide for use on Kentucky bluegrass grown for seed.

**This label for EVEREST 2.0 Herbicide expires and must not be distributed or used in accordance with this SLN registration after March 15, 2016.**

**ARYSTA LIFESCIENCE NORTH AMERICA, LLC  
15401 Weston Parkway, Suite 150  
Cary, North Carolina 27513**

EVEREST is a registered trademark of Arysta LifeScience North America, LLC.  
Label Revision Date: March 3, 2011