

syngenta.

INTIRA

Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF IDAHO

Fusilade® DX
Fine Fescue Grasses (Chewings, Hard, and Creeping Red Varieties) Grown
for Seed Use Only

EPA Reg. No. 100-1070

EPA SLN No. ID-040003

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2019

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.
- **FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

Fine Fescues Grown for Seed

For control of downy brome (cheatgrass) (*Bromus tectorum*), quackgrass (*Agropyron repens*), bentgrass (*Agrostis sp.*), and volunteer cereals in Fine Fescue Grasses (such as Creeping Red, Chewings and Hard Fescue varieties) grown for seed in Idaho. **NOT FOR USE ON TALL FESCUE.** Refer to the federal label for a complete listing of grass weeds controlled.

Fall Application - Apply Fusilade DX at 8 oz per acre to actively growing downy brome in the 2 to 4 inch stage. Apply at 12 to 16 oz per acre to actively growing quackgrass or bentgrass in the 6 to 10 inch stage. If regrowth of quackgrass or bentgrass occurs, make a second application at 12 to 16 oz per acre before regrowth exceeds 10 inches. **Spring Application** - Apply Fusilade DX at 8 to 16 oz per acre to actively growing grasses. Downy brome in the 2 to 4 inch stage can be controlled by the lower rate but larger, established annual grasses (4 to 6 inches) may need the higher rate for adequate control. . Apply at 12 to 16 oz per acre to actively growing quackgrass or bentgrass in the 6 to 10 inch stage. If regrowth of quackgrass or bentgrass occurs, make a second application at 12 to 16 oz per acre before regrowth exceeds 10 inches.

RESTRICTIONS

- Do not apply FUSILADE DX after fescue grass seedhead develops into the boot stage.
 - Do not apply this product through any type of irrigation system.
 - Treated fields may not be grazed or harvested for forage until after harvesting grass seed crop.
 - For use only on fields of fine fescue (chewings, creeping red, and hard fescues) grown for seed. Other types of fescue may not be tolerant.
 - Do not apply a total of more than 64 fl oz of Fusilade DX per acre per year to fine fescue grasses grown for seed.
-

Fusilade® and the Syngenta logo trademarks of a Syngenta Group Company

©2014 Syngenta

24(c) Registrant:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, NC 27419-8300

Label Code: ID1070062BA0414