

FIFRA Section 24(c) Special Local Need



GROUP	3	FUNGICIDE
-------	----------	-----------

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF IDAHO



FOR USE ON CANOLA THROUGH CHEMIGATION
 EPA Reg. No. 59639-147
 EPA SLN No. ID110005

Active Ingredient	By Wt.
*Metconazole	50%
Other Ingredients	50%
Total	100%

*5-[(4-chlorophenyl)methyl]-2,2-dimethyl-1-(1*H*-1,2,4-triazol-1-ylmethyl)cyclopentanol

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Persons using this product must comply with all applicable directions, restrictions, Worker Protections Standard requirements and precautions found on the label of the Federally registered product upon which this registration is based.

This labeling must be in the possession of the user at the time of pesticide application. **READ ENTIRE LABEL AND BOOKLET. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

APPLICATION DIRECTIONS		
Crop	Diseases Controlled	Application Rate
Canola	White Mold/Sclerotinia Stem Rot <i>(Sclerotinia sclerotiorum)</i>	2.0 to 4.0 oz/A (0.0625 to 0.125 bl ai/A)

APPLICATION THROUGH IRRIGATION SYSTEMS (CHEMIGATION) FOR CANOLA

Quash alone or in combination with other pesticides which are registered for application through irrigation systems, may be applied through irrigation systems. Apply *Quash* only through wheel lines, linear, center pivot and solid set irrigation. Do not apply through any other type of irrigation system. Illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have any questions about calibration, you should contact State Extension Service specialist, equipment manufacturers, or other experts. Do not connect an irrigation system used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

USE RESTRICTIONS:

1. Make application between 20% and 50% bloom.
2. Apply via chemigation using 1/8 to 1/3 inch of water over the area to be treated.
3. Under high disease pressure, use the application rate of 4 oz/A
4. Do not apply more than 4 oz/A per season.
5. Do not apply more than one application per season.
6. Do not apply within 35 days of harvest

FIFRA § 24(c) registrant: Valent U.S.A. Corporation
P.O. Box 8025
Walnut Creek, CA 94596-8025

Quash is a registered trademark of Valent U.S.A. Corporation

**THIS LABELING EXPIRES AFTER DECEMBER 31, 2015 AND MUST NOT BE USED OR
DISTRIBUTED AFTER THAT DATE.**